

Status of the HIRDLS HNO₃ Data Product

D. Kinnison, J. Gille, J. Barnett, C. Randall, L. Harvey, S. Massie, C. Halvorson, B. Nardi, A. Lambert, H. Lee, M. Coffey, T. Eden, R. G. Francis, C. Cavanaugh, C. Craig, J. McInerney, C. Krinsky, B. Peterson, J. Craft, V. Dean, J. Reburn, A. Waterfall, and C. Waymark

+ MLS Science Team

+ ACE Science Team

+ FIRS-2 Science Team

+ CIMS Science Team

Aura Science Team Meeting

11-15 September 2006




Presentation Outline




- HIRDLS HNO_3 Signal Characteristics
- HIRDLS HNO_3 Climatological Distributions
- HIRDLS HNO_3 Vertical Range
- HIRDLS HNO_3 Accuracy
 - Compared to ACE
 - Large Balloon (FIRS-2, 2005)
 - PAVE
 - Houston Ave 2005
- Summary and Future Algorithm Development
 - Validation paper?

HNO₃ Channel 8 Radiance Signal is Strong!

- Atmospheric radiances from HIRDLS channels span ~ 4 orders of magnitude
- Initial success for T and O₃ was for channels with largest radiances, most tolerant to correction errors



Limb Radiance Spectra for HIRDLS channels 6-9 for a Tangent Height of 25km



Edwards et al., Appl. Optics, 1995.

HNO₃ has a strong radiance signal in channel 8 between 861-903 cm^{-1}


Presentation Outline




- HIRDLS HNO_3 Signal Characteristics
- HIRDLS HNO_3 Climatological Distributions
- HIRDLS HNO_3 Vertical Range
- HIRDLS HNO_3 Accuracy
 - Compared to ACE
 - Large Balloon (FIRS-2, 2005)
 - PAVE
 - Houston Ave 2005
- Summary and Future Algorithm Development
 - Validation paper?

Global Comparisons January (1979; 2006)

binned: 5° latitude; 20° longitude; all profiles



Correct Winter/Summer Asymmetry represented in HIRDLS (more HNO₃ in NH)




Cloud signatures


Retrieval issues in the upper stratosphere, polar region

Global Comparisons May 11th (1979; 2006)

binned: 5° latitude; 20° longitude; all profiles




Correct Winter/Summer Asymmetry represented in HIRDLS (more HNO₃ in SH)



HIRDLS VMR peak too high.


HNO_3 (ppbv) *** Orbit Plot for 2005, September 20



HIRDLS V2.02 vs
MLS V2.1

Common Features



More noise in polar high altitude region.



Presentation Outline



- HIRDLS HNO_3 Signal Characteristics
- HIRDLS HNO_3 Climatological Distributions
- HIRDLS HNO_3 Vertical Range
- HIRDLS HNO_3 Accuracy
 - Compared to ACE
 - Large Balloon (FIRS-2, 2005)
 - PAVE
 - Houston Ave 2005
- Summary and Future Algorithm Development
 - Validation paper?





HIRDLS HNO₃ Profile.

Latitude = 31°S

Longitude = 12°W

- σ = Standard deviation of the total error from the diagonal of the error covariance matrices.
- Typically all profiles have a σ = < 20% between 100 hPa and 10 hPa.
- When the variance of the total error increases significantly greater 20-30%, the contribution of the *a priori* increases dramatically.



HIRDLS HNO₃ Profile.

Latitude = 57°N

Longitude = 15°W

Some profiles have a larger range: 6 hPa to 250 hPa.

Presentation Outline



- HIRDLS HNO_3 Signal Characteristics
- HIRDLS HNO_3 Climatological Distributions
- HIRDLS HNO_3 Vertical Range
- HIRDLS HNO_3 Accuracy
 - Compared to ACE
 - Large Balloon (FIRS-2, 2005)
 - PAVE
 - Houston Ave 2005
- Summary and Future Algorithm Development
 - Validation paper?

Atmospheric Chemistry Experiment



Launched on the Canadian SCISAT-1 satellite on 12 Aug 2003

IR (2.2 – 13.3 μm) Fourier Transform Spectrometer (0.02 cm^{-1} resolution)

Solar Occultation, 74° Inclination:

Near global coverage in ~ 1 month, but extended periods in polar region
Sunrise/Sunset only

Vertical resolution: ~ 4 km

Cloud Top to 150 km

(See *GRL special issue*, vol. 32, 2005.)

Courtesy of
Cora
Randall

HIRDLS vs. ACE Comparisons



HIRDLS data are version 2.02.

ACE data are near-real-time version 2.2.

- Do *not* use High altitude HNO_3 retrievals because there are not enough retrievals at this time.


Data Screening

- Omitted HIRDLS data poleward of 63°S .
- Omitted HIRDLS data if precision was negative.
- Omitted ACE data if too large a contribution from a priori.

Coincidence Criteria: Same Day, ± 500 km.

All **HIRDLS** profiles coincident with a single **ACE** profile were averaged together before comparing.

HIRDLS/ACE Coincidences in May 2006



Total of 409 coincidences

262 in NH

147 in SH

Avg Separation = 170 km,
ranging from 7-470 km

NH ACE:

4-22 May Local SR

25-31 May Local SS

NH HIRDLS:

4-22 May ~5 hrs before
to 20 hrs after ACE

25-31 May 5-15 hrs
before ACE


SH ACE:

Local SS for all
coincidences

SH HIRDLS:

~5 hours later than ACE

HIRDLS/ACE Coincidences in May 2006




Similar to map on previous slide, but this shows the average location for all HIRDLS profiles (often ~10) coincident with a single ACE profile.

HIRDLS vs ACE




HIRDLS (red) vs. ACE (blue) HNO₃


NH HNO₃




SH HNO₃



NH HNO₃



SH HNO₃




Courtesy of
Cora
Randall


HIRDLS vs ACE




NH HNO₃




NH HNO₃ (%)



SH HNO₃



SH HNO₃ (%)




Courtesy of
Cora
Randall

Coincidence Criteria: ± 2 hrs, ± 500 km.

All **HIRDLS profiles** coincident with a single **ACE profile** were averaged together before comparing.

HIRDLS/ACE Coincidences, 18-31 May & 11-13 Jul, 2006



Total of 156
coincidences

18-31 May 2006

11-13 July 2006





113 in NH
43 in SH

Avg Separation =
302 km, ranging
from 37-495 km

Avg Time Diff =
1.1 hrs

Courtesy of
Cora
Randall

HIRDLS & ACE Nitric Acid Profiles




All
Coincidences

Average
(solid) & 1- σ
standard
deviation
(dotted)


Courtesy of
Cora
Randall

HIRDLS-ACE Nitric Acid Differences


NH HNO₃




NH HNO₃ (%)



SH HNO₃



SH HNO₃ (%)



Thick red:
Average

Dotted red:
1-σ distribution

Thin red:
1-σ uncertainty
(often hidden)

Black points:
Individual
differences

Courtesy of
Cora
Randall

Far-Infrared Spectrometer (FIRS-2)



Launched in New Mexico, Ft. Sumners.

High Resolution Fourier Transform Spectrometer (0.004 cm^{-1} resolution)


Measures thermal emissions of atmosphere from balloon.

Vertical resolution: 3-4 km

Cloud Top to approx 40km (3 hPa)

(See *Jucks et al., 1998; ...*)


HNO₃ *** 31.4N, 254.9E, *** 20050920



HIRDLS vs FIRS-2

- September 20, 2005
 - 31.4 N latitude
 - 255 E longitude.
- Coincidence is within 5° longitude and 2° latitude.
- 8 HIRDLS profiles
- Daytime overpasses for HIRDLS
- Used the FIRS-2 profile that was nearest to the Aura overpass.
- Preliminary FIRS-2 data (Ken Jucks).


HIRDLS vs NCAR FTS (See M. Coffey's Poster)



- PAVE February 7, 2005
- Comparisons are made outside of the vortex on this day.

Courtesy of Mike Coffey


HIRDLS vs MLS and NCAR FTS



- PAVE February 7, 2005
- MLS V1.5

Courtesy of Mike Coffey


HIRDLS vs NCAR FTS (See M. Coffey's Poster)



- PAVE February 5, 2005
- MLS V2.1 data

Courtesy of Mike Coffey


HIRDLS vs MLS and NCAR FTS




- PAVE February 5, 2005
- MLS V2.1 data

Courtesy of Mike Coffey

[2] 2005/06/19 flight track superimposed on GOES-12 IR image



HNO₃ *** AVE Houston *** 20050619



HIRDLS vs CIMS

- Houston AVE, June 19th 2005.
- HIRDLS profiles between 28-34°N (8-total).
- CIMS data during descent. Downward facing inlet.
- HIRDLS agrees with CIMS within a factor of two.

Presentation Outline



- HIRDLS HNO_3 Signal Characteristics
- HIRDLS HNO_3 Climatological Distributions
- HIRDLS HNO_3 Vertical Range
- HIRDLS HNO_3 Accuracy
 - Compared to ACE
 - Large Balloon (FIRS-2, 2005)
 - PAVE
 - Houston Ave 2005
- Summary and Future Algorithm Development
 - Validation paper?

HNO₃

Summary...

- Morphology consistent with MLS and ACE
 - Orbit plot comparison between HIRDLS and MLS show similar structure.
 - High latitudes have correct hemispherical asymmetry.
 - Peak HNO₃ is (near) the right altitude.
- Vertical range of HNO₃ product is between 100 hPa 10 hPa
 - In some instances the range can be extended downward to approximately 200 hPa and upward to 3 hPa.
- HIRDLS Biased low relative to ACE, FIRS-2, FTS, CIMS
 - Biased low in the middle stratosphere (50%, 2-3 ppbv)
 - Biased high in the upper stratosphere (50%, 1-2 ppbv)
 - Larger bias in the SH vs NH.
 - Within a factor of two in the UTLS region.
- Overall HNO₃ product has improved greatly since September 2005.

The END